

## PRIMER NOTE

# Characterization of 35 microsatellite loci in the Pacific lion-paw scallop (*Nodipecten subnodosus*) and their cross-species amplification in four other scallops of the Pectinidae family

ANA M. IBARRA,\* JESSICA L. PETERSEN,† THOMAS R. FAMULA† and BERNIE MAY†

\*Aquaculture Genetics Laboratory, Centro de Investigaciones Biológicas del Noroeste S.C., A.P. 128, La Paz B.C.S., Mexico 23000,

†Department of Animal Science, University of California — Davis, 1 Shields Ave., Davis CA 95616, USA

## Abstract

Four microsatellite-enriched DNA libraries yielded 35 microsatellite loci from 100 primer pairs designed for Pacific lion-paw scallop, *Nodipecten subnodosus*. The number of alleles ranged from four to 28. Three of the 35 loci were not in Hardy–Weinberg equilibrium and linkage disequilibrium was found for one pair of loci. These microsatellites will be used to analyse the population structure of the species in Mexico's Baja Peninsula to propose management strategies for scallop aquaculture development. Twenty-six primer pairs cross-amplified in *Nodipecten nodosus*, whereas none (*Argopecten ventricosus*) or few cross-amplified in the *Argopecten* species.

**Keywords:** *Argopecten*, Guerrero Negro — Ojo de Liebre lagoon, microsatellite, primers

Received 13 July 2005; revision accepted 25 August 2005

The lion-paw scallop, *Nodipecten (Lyropecten) subnodosus*, is the largest scallop species found in the tropical waters of the East Pacific (Peña 2001). In Mexico, it is harvested on the Pacific and Gulf of California sides of the Baja California Peninsula and on the coast of Sonora state. The largest population is found in the lagoon of Ojo de Liebre, which is part of the Guerrero Negro lagoon system on the Pacific side of the Baja Peninsula. A fundamental question for the optimal development of lion-paw scallop aquaculture is whether populations native to different lagoons are genetically distinct. We isolated and characterized microsatellite loci that will allow for the analysis of population structure, from which optimum management strategies for aquaculture ventures can be proposed. These markers will also allow for evaluation of optimum spawning strategies for hatcheries that minimize genetic drift and inbreeding in this functional hermaphrodite species.

Whole genomic DNA was extracted using QIAGEN's DNeasy Tissue Kit from adductor muscle of wild lion-paw scallops naturally settled at Bahia de La Paz. DNA

obtained from four scallops and 18 crayfish, the last also being studied in the Genomic Variation Laboratory (GVL), were mixed in equal concentrations prior to library construction. Four libraries enriched for tetra- and trinucleotide repeat motifs (TAGA)<sub>n</sub>, (TGAC)<sub>n</sub>, (TACA)<sub>n</sub> and (ATC)<sub>n</sub> were constructed, screened and sequenced ( $n = 170$ ) from this mixed DNA by Genetic Identification Services (Chatsworth) according to Meredith & May (2002) and Schwartz & May (2004). One hundred and four primers were designed using PRIMERSELECT 4.0 (DNA Star Inc.) at the GVL of University of California — Davis.

To determine if loci amplified in scallop or crayfish, primers were first tested on four scallops from the same wild population. All 35 'positive' primers for the scallop were screened on 30 individuals, including 25 from Ojo de Liebre lagoon on the Pacific side of the Baja California Peninsula, and five from Bahia de La Paz, on the Gulf of California side of the peninsula. We further tested these primers for cross-amplification of the microsatellite loci in five to six individuals of each of the following scallop species: *Argopecten irradians* (USA, west Atlantic), *Argopecten purpuratus* (Chile, east Pacific), *Argopecten ventricosus* (Mexico, east Pacific) and *Nodipecten nodosus* (Venezuela, west Atlantic). DNA was extracted from muscle tissue

Correspondence: A. M. Ibarra, Fax: 011-52-612-125-3625; E-mail: aibarra@cibnor.mx

preserved in 70% alcohol (*A. irradians* and *A. ventricosus*), or from dried adductor muscle tissue (*A. purpuratus* and *N. nodosus*).

Polymerase chain reaction (PCR) was performed using 5 ng of genomic DNA, 1 × *Taq* DNA polymerase buffer B, 2.0 mM MgCl<sub>2</sub>, 0.2 mM of each dNTP, 1 μM of each primer and 0.38 U *Taq* DNA polymerase (all reagents from Promega), with a total PCR volume of 10 μL. PCR was carried out using an MJ Research PTC-100 under the following conditions: 94 °C for 2.30 min, 30 cycles at 94 °C for 30 s, 56 °C for 30 s (53 °C for cross-amplifications), 72 °C for 30 s, followed by 72 °C for 5 min, and held at 4 °C. For those primers producing multiple-bands in the primary species (*N. subnodosus*), we performed touchdown PCR: 95 °C for 1 min followed by 30 cycles at 95 °C for 1 min, 67 °C for 45 s with a 0.5 °C decrease each cycle, and 72 °C for 2 min, using the same reagents and equipment. The amplified products were diluted 1:1 with 98% formamide loading buffer, denatured at 95 °C for 2 min and chilled on ice for 2–3 min. Samples were separated on a 5% denaturing polyacrylamide gel at 50 W for 70 min. Amplified products were visualized using the Sybr-Green™-agarose overlay protocol (Rodzen *et al.* 1998) and scanned with a Molecular Dynamics FluorImager 595. Product sizes were estimated by comparison with a standard 400 bp ladder (The Gel Company).

The primer sequences amplifying in *N. subnodosus*, their polymorphism data and analyses for Hardy–Weinberg equilibrium (HWE) and linkage disequilibrium (LD), are presented in Table 1. The repeat motifs are specific to the one sequenced clone for each locus. Observed and expected heterozygosities were calculated from the 25 individuals from Ojo de Liebre lagoon. For the five individuals from Bahia La Paz, we present number of alleles and their size ranges. Deviations from HWE and LD were calculated using GDA program (Lewis & Zaykin 2001), and significance was evaluated with the Fisher’s exact test. Three loci were not in HWE after applying a Bonferroni correction, *NsubA007*, *NsubA010* and *NsubB215*, although it is not known if these deviations are due to null alleles or nonrandom mating because of the self-fertilizing capability for this functional hermaphrodite. Significant pairwise LD was found only between loci *NsubA223* and *NsubC262* following a test using genotypes to prevent within locus disequilibrium from affecting the significance and a Bonferroni correction for multiple comparisons ( $P \leq 0.0014$ ).

Results of cross-amplification of primers on the other four species are in Table 2. Of the 35 primer pairs developed, nine (25%) resulted in no amplification for all the cross-amplified species. None of the primers amplified in *A. ventricosus*. Twenty-six (74%) of the 35 primer pairs that successfully amplified polymorphic loci in *N. subnodosus* amplified in *N. nodosus*, the Atlantic species from the same genus. The next largest cross-amplification was seen for *A. irradians* (11 primer pairs) and for *A. purpuratus* (seven primer pairs).

**Table 1** Characterization of 35 microsatellite loci in lion-paw scallop (*Nodipecten subnodosus*) from Lagoon Ojo de Liebre (OL) within the Guerrero Negro lagoons system and Bahia de La Paz, Mexico. GenBank Accession nos, primer sequences, repeat motif, loci amplified using touchdown PCR, number of individuals genotyped, number of alleles, get estimated allele size range (bp), observed and expected heterozygosities, and test for conformance to Hardy–Weinberg equilibrium for the Ojo de Liebre lagoon. Allele number and size ranges for Bahia de La Paz

Locus	GenBank Accession no.	Primer sequence (5'–3')	TD PCR	n	Repeat motif	No. of alleles	Get estimated allele size range (bp)	H <sub>O</sub>	H <sub>E</sub>	P(HWE) & significance <sup>(1)</sup>	Bahia de La Paz (n = 5) no. of alleles (size range)
<i>NsubA001</i>	DQ108619	F: GAGGATGACAGTGTGAAGATG R: CACAAAACAGAAATGTGAAGA	+	23	(TGA) <sub>3</sub> TTA(TGA) <sub>9</sub>	4	285–300	0.65	0.54	0.6166	4 (285–300)
<i>NsubA004</i>	DQ108620	F: TTTGACAGACATCCCTTCTTAC R: GAATGCGACATAGACATCG	+	21	(TCA) <sub>10</sub>	9	230–267	0.86	0.86	0.7844	4 (237–247)
<i>NsubA005</i>	DQ108621	F: CGACAAACATCCCTCTTT R: ACAGCGACCAGTGTAAAG	+	25	(TCA) <sub>12</sub> AAATCA	4	204–218	0.80	0.66	0.2937	5 (204–218)
<i>NsubA007</i>	DQ108622	F: AACGCTGTCACTGAAACAAAG R: TCGGTGATACAAAGTTGAAGGG	+	21	(CAT) <sub>18</sub> (GTC) <sub>5</sub> CAT(CGT) <sub>4</sub> (C) <sub>11</sub>	13	159–285	0.48	0.91	0.0000*	5 (200–242)
<i>NsubA010</i>	DQ108623	F: TTTGACTTGAAGACATCTCT R: CGGCACATCAITGTTACTAAA	+	22	(TCA) <sub>2</sub> N <sub>10</sub> TCAATCATACA(TCA) <sub>3</sub>	6	126–144	0.95	0.74	0.0013*	4 (128–138)
<i>NsubA208</i>	DQ108624	F: GTTACGTGGTGGACTGAA R: CGGCATCTGTACTGACAC	+	24	(ATG) <sub>9</sub>	4	283–298	0.58	0.54	0.8284	4 (287–298)
<i>NsubA214</i>	DQ108625	F: CCTCTCTCCACTTCTTTTTCAC R: CACCCGGAGCTTAAATATPAGG		22	(TCA) <sub>10</sub> TA(TCA) <sub>11</sub>	5	235–290	0.50	0.67	0.0069	3 (235–278)
<i>NsubA222</i>	DQ108626	F: TTTTGTGACGCCAACACTA R: GACAGGGACCGACTTACC		22	(TCA) <sub>21</sub> (TCG) <sub>5</sub> (TCA) <sub>2</sub> TCG(TCA) <sub>3</sub> (TCG) <sub>9</sub>	9	220–320	0.91	0.80	0.6237	5 (220–300)
<i>NsubA223</i>	DQ108627	F: CCTGTCTTTTGAACGTCTAG R: CTGTGCTGTCTTCAGACAAAC		24	(CAT) <sub>5</sub> TAAT(CAT) <sub>6</sub> CAA(TCA) <sub>15</sub>	14	120–240	0.79	0.90	0.1053	8 (120–200)
<i>NsubA227</i>	DQ108628	F: ACCTCATCTGGTGTCAATTC R: CAATTTCCAGTGTGTGTGAT		21	(ATC) <sub>21</sub> (TCG) <sub>12</sub>	13	180–300	0.95	0.92	0.6134	5 (180–300)

Table 1 Continued

Locus	GenBank Accession no.	Primer sequence (5'–3')	TD PCR	<i>n</i>	Repeat motif	No. of alleles	Gel estimated allele size range (bp)	$H_O$	$H_E$	P(HWE) & significance <sup>(1)</sup>	Bahía de La Paz ( <i>n</i> = 5) no. of alleles (size range)
<i>NsubA229</i>	DQ108629	F: TCTCCCTGGAATGATAAAGG R: GGCTGTAAATCTCACCAATG		19	(ACG) <sub>5</sub> N <sub>9</sub> (ATG) <sub>4</sub> A(CGA) <sub>8</sub> (TGA) <sub>6</sub>	7	242–284	1.00	0.83	0.3769	4 (242–280)
<i>NsubA231</i>	DQ108630	F: CTGGGAATTTGTAGCTTTTACC R: GAGGAAGATGATGATGATGATT	+	25	(TCA) <sub>5</sub> TCTCTTAAA (TCA) <sub>4</sub>	4	138–150	0.48	0.57	0.1647	3 (138–147)
<i>NsubA235</i>	DQ108631	F: GTGGCAAGTCTTATTTGGTAGA R: CTGTTTACTTTTCGATGCAAGTT		24	(TGA) <sub>11</sub>	7	200–230	0.71	0.78	0.4853	5 (200–225)
<i>NsubA238</i>	DQ108632	F: TTTTATCGAAATGTCGGATGTC R: TGCTGGTCGAGTTCTTAAAGAG		24	(TCA) <sub>4</sub> N <sub>7</sub> (TCA) <sub>8</sub> N <sub>6</sub> (TCC) <sub>8</sub>	13	98–155	1.00	0.92	0.5319	6 (110–155)
<i>NsubA243</i>	DQ108633	F: CGCAGGAAACGATCATTTAGTA R: CACACGGAACATCAACTTAACC	+	25	(GAT) <sub>7</sub> AT(GAT) <sub>11</sub>	7	142–165	0.88	0.78	0.0078	5 (142–158)
<i>NsubA245</i>	DQ108634	F: TGCCATTTTCGATAACTGATG R: TAATGCTAATACGGCACCAGT		25	(TCA) <sub>8</sub>	6	293–320	0.88	0.81	0.2313	4 (305–320)
<i>NsubA249</i>	DQ108635	F: CCGCTGAAAATCTCTCCT R: GCCCATCGTAAACAATCT	+	21	(TCA) <sub>13</sub>	8	240–295	1.00	0.85	0.1025	6 (240–278)
<i>NsubA261</i>	DQ108636	F: ACGAAAAGTTGAAATCGTGATC R: CGTTTATACAGGCAATGTTCTG	+	22	(CAT) <sub>6</sub> T(ATC) <sub>6</sub> (GTC) <sub>11</sub>	9	240–320	0.95	0.87	0.6572	4 (240–280)
<i>NsubA262</i>	DQ108637	F: ATCACGAGTCAAAAACAGTATCG R: CCTGGTAGGTGAATCACACTAA		20	(ATC) <sub>4</sub> N <sub>18</sub> (ATC) <sub>8</sub>	12	285–430	0.70	0.89	0.0084	6 (290–405)
<i>NsubA266</i>	DQ108638	F: GCCAAATTTATGACATCCAC R: AATGGGTTGATGATGATGTC	+	21	(TCA) <sub>9</sub> G(CAT) <sub>3</sub>	14	132–225	0.90	0.89	0.5334	6 (147–210)
<i>NsubA274</i>	DQ108639	F: GAACTTGGTGCAGTTATCTTCA R: ATAGCGTAAAGTGTGCAACGT	+	22	(TCA) <sub>9</sub>	12	260–325	0.95	0.87	0.5650	6 (262–290)
<i>NsubB007</i>	DQ108640	F: GGCATGTGAACACTACATTTCAAT R: TTCCATTCGGTTTACACAGATA		22	(ATGT) <sub>12</sub>	8	260–320	0.77	0.83	0.5241	5 (270–295)
<i>NsubB210</i>	DQ108641	F: GAGAGGTGAAGTGAAGTGAAG R: TGGAAGTACAGAAAGGAATGTC		20	(CATA) <sub>12</sub>	5	240–290	0.60	0.76	0.0016	5 (240–290)
<i>NsubB215</i>	DQ108642	F: CTGACTTATCCTGTCATTTCA R: TATTCCTTTCTCGCCCATAT		25	(ATAC) <sub>13</sub> N <sub>6</sub> (ATAC) <sub>4</sub> N <sub>6</sub> (ATAC) <sub>15</sub>	28	95–420	0.80	0.97	0.0000*	9 (95–400)
<i>NsubB235</i>	DQ108643	F: GTCCTTTGGTCTTGGTTACTG R: ATGGATTTGACATCATCAAC		22	(TACA) <sub>4</sub> CA(TACA) <sub>5</sub> TACG(TACA) <sub>3</sub>	5	145–180	0.86	0.75	0.3788	5 (145–180)
<i>NsubB252</i>	DQ108644	F: ACAAGCGAATACAACGACA R: ACCATTGACCAGGCTACAT	+	22	(TGTA) <sub>4</sub> TGGTA(TGTA) <sub>9</sub>	4	150–178	0.73	0.72	0.5375	4 (150–178)
<i>NsubB278</i>	DQ108645	F: CACACGATTCACCTTTCCTATTG R: GCTTCAGCTTAACGATTTGAAAC	+	19	(TACA) <sub>12</sub> CA(TACA) <sub>13</sub> CA(TACA) <sub>11</sub>	17	138–440	1.00	0.93	0.9794	6 (142–358)
<i>NsubC020</i>	DQ108646	F: CACCTACCTCTAGGTGTGGAAG R: TGTGAAAGACAGAGTTATGATCGAG		21	(TGTC) <sub>4</sub> ATGTCGTGACC (TATC) <sub>7</sub>	10	195–280	0.81	0.88	0.2491	8 (195–270)
<i>NsubC023</i>	DQ108647	F: CCCCAAAACACAATGCAGAACAAA R: TTATGGCCGAAAGTATCAATCAG		23	(TAGA) <sub>16</sub> (TGGG) <sub>2</sub> TGA (TGGG) <sub>8</sub> (TAGA) <sub>2</sub> (AGA) <sub>16</sub>	20	250–520	0.96	0.96	0.5822	4 (360–500)
<i>NsubC205</i>	DQ108648	F: GTCAAAAGGTGGCAGTCTAAG R: CAGCATAAACCCCTACTCATGGT	+	21	(GATA) <sub>10</sub>	9	195–240	0.95	0.84	0.6253	2 (195–202)
<i>NsubC218</i>	DQ108649	F: CCGCAAATATCATTTATCTCG R: GTCGTTAAAGGTTGGTGTG	+	18	(GTCT) <sub>4</sub> (ATCT) <sub>14</sub>	9	170–320	0.94	0.89	0.4478	7 (170–290)
<i>NsubC250</i>	DQ108650	F: ACCTGAAGCAAAGAAAGAAAGA R: TTACCTGTAGGTGGGAGACAC		25	(GACA) <sub>4</sub> (GATA) <sub>4</sub> CATA(GATA) <sub>4</sub>	11	250–320	0.84	0.89	0.8634	8 (255–300)
<i>NsubC261</i>	DQ108651	F: TTCAAACGCACGTCATTTG R: CTTGGCACCATATTCGGAC		24	(TATC) <sub>8</sub>	9	125–200	0.71	0.85	0.0031	4 (145–160)
<i>NsubC262</i>	DQ108652	F: CCGCAAATATCATTTATCTCG R: GCATTGTCAACTGAATTAGCA		23	(TGTC) <sub>4</sub> (TATC) <sub>10</sub>	15	155–250	0.74	0.93	0.1219	4 (168–217)
<i>NsubC275</i> <sup>(2)</sup>	DQ108653	F: AGGCTGGGTCACATATCA R: CGGTGATTACAAAATACACC	+	22	(TATC) <sub>7</sub>	5	78–114	0.86	0.72	0.5216	3 (78–90)

+ Best amplification – resolution by using TD PCR.

\*Statistically significant ( $P < 0.05$ ).

(1) After standard Bonferroni correction.

(2) Forward primer within minisatellite; amplification at 3 other size ranges (c. 130–150, 200–220 &amp; 250–270).

**Table 2** Cross-species amplification results (number of alleles per locus with size range (bp)) in parentheses of 35 microsatellite loci for the scallop family Pectinidae, genus *Argopecten* (*A. irradians*, *A. purpuratus*, *A. ventricosus*) and *Nodipecten* (*N. nodosus*)

Locus ID	<i>A. irradians</i> (n = 6) USA — Atlantic	<i>A. purpuratus</i> (n = 5) Chile — Pacific	<i>A. ventricosus</i> (n = 6) Mexico — Pacific	<i>N. nodosus</i> (n = 6) Venezuela — Atlantic
<i>NsubA001</i>	U	—	—	U (140, 215)
<i>NsubA004</i>	—	—	—	2 (215–235)
<i>NsubA005</i>	—	—	—	3 (180–200)
<i>NsubA007</i>	U	U	—	6 (160–220)
<i>NsubA010</i>	U (115, 300)	—	—	4 (120–135)
<i>NsubA208</i>	—	—	—	3 (280–300)
<i>NsubA214</i>	—	—	—	—
<i>NsubA222</i>	—	—	—	—
<i>NsubA223</i>	U	—	—	7 (90–120)
<i>NsubA227</i>	—	—	—	U (200–240)
<i>NsubA229</i>	—	—	—	—
<i>NsubA231</i>	4 (300–400)	U (340, 460)	—	5 (140–150)
<i>NsubA235</i>	—	6 (200–300)	—	7 (200–235)
<i>NsubA238</i>	—	—	—	U (120–140)
<i>NsubA243</i>	—	—	—	U (120–160)
<i>NsubA245</i>	—	—	—	—
<i>NsubA249</i>	—	—	—	2 (250–270)
<i>NsubA261</i>	—	—	—	7 (220–300)
<i>NsubA262</i>	—	U	—	5 (280–340)
<i>NsubA266</i>	—	—	—	—
<i>NsubA274</i>	U	—	—	3 (260–280)
<i>NsubB007</i>	—	—	—	—
<i>NsubB210</i>	—	—	—	—
<i>NsubB215</i>	—	—	—	5 (100–300)
<i>NsubB235</i>	—	—	—	U
<i>NsubB252</i>	U	—	—	2 (140–160)
<i>NsubB278</i>	—	—	—	—
<i>NsubC020</i>	—	—	—	2 (160–200)
<i>NsubC023</i>	—	—	—	—
<i>NsubC205</i>	2 (> 400)	U	—	3 (170–210)
<i>NsubC218</i>	U	U	—	4 (290–320)
<i>NsubC250</i>	2 (300–340)	—	—	4 (210–240)
<i>NsubC261</i>	—	—	—	3 (140–240)
<i>NsubC262</i>	—	—	—	6 (150–250)
<i>NsubC275</i>	U	U	—	U
Total no. of amplified loci	11	7	0	26

U, indicates amplification but unclear; '—' indicates no amplification; (num-num) indicates range; (num, num) indicates when two bp ranges amplified.

Species, sample size (n), and origin — distribution.

## Acknowledgements

UCMEXUS-CONACYT supported this research through grant (CN-04–61; B May — AM Ibarra) and a sabbatical fellowship (AM Ibarra). Samples from other species were donated by Amy Wilbur (*Argopecten irradians*), Iker Uriarte (*Argopecten purpuratus*) and Cesar Lodeiros (*Nodipecten nodosus*). Melinda Baerwald and Rick Topinka reviewed and commented on the manuscript.

## References

- Lewis PO, Zaykin D (2001) *GENETIC DATA ANALYSIS: computer program for the analysis of allelic data*, version 1.1. Free program distributed by the authors over the Internet from <http://hydrodictyon.eeb.uconn.edu/people/plewis/software.php> (Accessed July 2005).
- Meredith EP, May B (2002) Microsatellite loci in Lahontan tui chub, *Gila bicolor obesa*, and their utilization in other chub species. *Molecular Ecology Notes*, **2**, 156–158.
- Peña JB (2001) Taxonomía, morfología, distribución y hábitat de los pectínidos iberoamericanos. In: *Los Moluscos Pectínidos de Iberoamérica: Ciencia y Acuicultura* (ed. Maeda-Martínez AN), Chap. 1: pp. 1–25. Ed. Limusa, Mexico.
- Rodzen JR, Agresti JJ, Tranah GJ, May BP (1989) Agarose overlays allow simplified staining of polyacrylamide gels. *BioTechniques*, **25**, 584.
- Schwartz RS, May B (2004) Characterization of microsatellite loci in Sacramento perch (*Archoplites interruptus*). *Molecular Ecology Notes*, **4**, 694–697.